

# Dysphagia Screening Tool – Stroke Narrator ED

When caring for a Stroke patient, it is important to add the Dysphagia Screening Tool

1 This tool is available in the same location as the NIH scale

The screenshot shows the 'Past Stroke' patient record interface. The 'Dysphagia Screening' tool is highlighted in the right-hand navigation pane under 'Stroke Documentation'. The main window displays the tool's configuration, including 'Time taken: 08:09:08' on '4/11/2019'. The tool is set to show 'Row Info', 'Last Filed', and 'All Choices'. The 'Swallow Function' section is expanded, showing various symptoms like 'continuously clears throat', 'cough', 'difficulty initiating swallow', and 'difficulty tolerating secretions'. Other sections include 'Reason for deferral', 'Follow up', and 'Deferral'.

2 Follow the instructions listed in the group and row information

This screenshot provides a closer view of the 'Dysphagia Screening' tool. The 'Time taken' is now '08:08' on '4/11/2019'. The 'Swallow Function' section is expanded, showing a list of symptoms: 'continuously clears throat', 'cough', 'difficulty initiating swallow', 'difficulty tolerating secretions', 'drools', 'excessive chewing', 'frequently washes down food with liquid', 'moves head/neck while swallowing', 'plays with food', 'pockets food', 'repeated swallowing of single bite', 'spills food from mouth', 'swallows without difficulty', 'throat discomfort while swallowing', 'voice quality change', and 'other (see comments)'. Below this, there are sections for 'Reason for deferral', 'Follow up', and 'Deferral'. The 'Deferral' section includes options like 'Due to presenting complaint', 'Unable to assess - infant', and 'Unable to assess - trauma'. The 'Accept' and 'Cancel' buttons are visible at the bottom.